

# A New Class of Bipyridine-ligated Metal Carboxylate Complexes: Characterization of the Triply-bridged Ferromagnetically-coupled Complexes $[\text{Cu}_2(\text{OAc})_3(\text{bpy})_2](\text{ClO}_4)$ and $[\text{Cu}_2(\text{OH})(\text{H}_2\text{O})(\text{OAc})(\text{bpy})_2](\text{ClO}_4)_2$

George Christou,<sup>a</sup> Spiros P. Perlepes,<sup>a</sup> Kirsten Folting,<sup>a</sup> John C. Huffman,<sup>a</sup> Robert J. Webb,<sup>b</sup> and David N. Hendrickson<sup>\*b</sup>

<sup>a</sup> Department of Chemistry and the Molecular Structure Centre, Indiana University, Bloomington, IN 47405, USA

<sup>b</sup> Department of Chemistry, University of California, San Diego, CA 92093, USA

Treatment of  $\text{Cu}_2(\text{OAc})_4 \cdot 2\text{H}_2\text{O}$  with 2,2'-bipyridine (bpy) and  $\text{NaClO}_4$  in EtOH yields  $[\text{Cu}_2(\text{OAc})_3(\text{bpy})_2](\text{ClO}_4)$  (**1**) whereas the same reaction in hot water yields  $[\text{Cu}_2(\text{OH})(\text{H}_2\text{O})(\text{OAc})(\text{bpy})_2](\text{ClO}_4)_2$  (**2**); the novel structures of the complexes are described, together with the results of magnetic susceptibility studies which show that both complexes possess intramolecular ferromagnetic coupling.


In recent years, we have been reporting our progress towards developing 3d transition metal carboxylate chemistry employing 2,2'-bipyridine (bpy) as a terminal ligand, seeking models for the various Mn biomolecules being identified.<sup>1</sup> Most of our efforts have hitherto been concentrated in Mn, and a growing family of complexes of various nuclearities has been obtained.<sup>1</sup> We recently reported preliminary extension of our work to Fe.<sup>2</sup> We have now extended our efforts still further to Cu, and we wish to report our first successes in this area, the preparation of two new dinuclear  $\text{Cu}^{\text{II}}$  complexes with novel bridging units.

Treatment of a stirred slurry of  $\text{Cu}_2(\text{OAc})_4 \cdot 2\text{H}_2\text{O}$  (1.00 g, 2.5 mmol) in EtOH (70 cm<sup>3</sup>) with solid bpy (0.78 g, 5.0 mmol) gave a clear blue solution. Addition of  $\text{NaClO}_4$  (0.61 g, 5.0 mmol) in EtOH (10 cm<sup>3</sup>) rapidly precipitated sky-blue microcrystalline  $[\text{Cu}_2(\text{OAc})_3(\text{bpy})_2](\text{ClO}_4)$  (**1**) in 95% yield that can be recrystallized from hot MeCN or MeCN/Et<sub>2</sub>O.<sup>†</sup>

In a similar fashion, a slurry of  $\text{Cu}_2(\text{OAc})_4 \cdot 2\text{H}_2\text{O}$  (1.00 g, 2.5 mmol) and  $\text{NaClO}_4$  (0.61 g, 5.0 mmol) in H<sub>2</sub>O (65 cm<sup>3</sup>) was treated with bpy (0.78 g, 5.0 mmol) dissolved in EtOH (10 cm<sup>3</sup>). The resulting slurry was heated to 75 °C and all solids dissolved to give a deep-blue solution. The solution was left to cool slowly overnight to ambient temperature to produce well-formed dark-blue crystals of  $[\text{Cu}_2(\text{OH})(\text{H}_2\text{O})(\text{OAc})(\text{bpy})_2](\text{ClO}_4)_2$  (**2**) in 69% yield. The product can be recrystallized from hot water.<sup>†</sup>

The structures<sup>‡</sup> of the two cations are shown in Figures 1 and 2. The structure of (**1**) consists of two  $\text{Cu}^{\text{II}}$  centres bridged by three  $\text{AcO}^-$  groups, two in the familiar  $\eta^1 : \eta^1 : \mu_2$  mode and the third in the rarer mono-atomic bridging mode,<sup>3-5</sup> with C(36)–O(35) [1.319(4) Å] significantly longer than C(36)–


O(37) [1.224(4) Å] as a result. A terminal bpy group completes five-co-ordination at each metal. As is sometimes<sup>4</sup> (but not always<sup>5</sup>) observed for mono-atomic bridging  $\text{RCO}_2^-$  groups, there is evidence for an additional weak interaction between Cu(1) and O(37). The Cu(1)···O(37) distance is only 2.716(2) Å, and the  $\text{AcO}^-$  group is clearly tilted towards Cu(1) as manifested in the Cu(1)–O(35)–C(36) angle [108.4(2)°] being significantly smaller than Cu(2)–O(35)–C(36) [127.3(2)°]. Also, the Cu(1)–O(35) bridging distance [1.977(2) Å] is noticeably shorter than the Cu(2)–O(35) distance [2.169(2) Å]. Further, the co-ordination geometries of the two Cu atoms are significantly different, and we assign this as also due to the Cu(1)···O(37) interaction. Thus, Cu(2) is intermediate between square pyramidal and trigonal bipyramidal whereas Cu(1) is approximately square pyramidal with O(31) at the apex. Indeed, this places O(37) approximately at the vacant sixth position of the Cu(1) square pyramid [O(31)–Cu(1)–O(37), 149.4(2)°].



**Figure 1.** The structure of the cation of complex (**1**). Selected distances (Å) and angles (deg). Cu(1)···Cu(2), 3.392(1); Cu(1)–O(27), 1.939(2); Cu(1)–O(31), 2.238(2); Cu(1)–O(35), 1.977(2); Cu(1)–N(3), 2.001(2); Cu(1)–N(14), 2.019(2); Cu(2)–O(29), 1.930(2); Cu(2)–O(33), 1.974(2); Cu(2)–O(35), 2.169(2); Cu(2)–N(15), 1.992(2); Cu(2)–N(26), 2.033(2); C(36)–O(35), 1.319(4); C(36)–O(37), 1.224(4); C(32)–O(31), 1.245(3); C(32)–O(33), 1.270(3); C(28)–O(27), 1.255(3); C(28)–O(29), 1.259(4); O(27)–Cu(1)–O(31), 97.4(1); O(27)–Cu(1)–O(35), 93.6(1); O(31)–Cu(1)–O(35), 96.5(1); O(27)–Cu(1)–N(3), 168.8(1); O(27)–Cu(1)–N(14), 88.3(1); O(31)–Cu(1)–N(3), 87.6(1); O(31)–Cu(1)–N(14), 102.4(1); O(31)–Cu(1)–O(37), 149.4(2); O(35)–Cu(1)–N(3), 95.9(1); O(35)–Cu(1)–N(14), 160.6(1); N(3)–Cu(1)–N(14), 80.8(1); Cu(1)–O(35)–C(36), 108.4(2); Cu(1)–O(35)–Cu(2), 109.8(1); O(29)–Cu(2)–O(33), 94.9(1); O(29)–Cu(2)–O(35), 96.0(1); O(29)–Cu(2)–N(15), 168.3(1); O(29)–Cu(2)–N(26), 88.6(1); O(33)–Cu(2)–O(35), 96.7(1); O(33)–Cu(2)–N(15), 90.6(1); O(33)–Cu(2)–N(26), 140.1(1); O(35)–Cu(2)–N(15), 93.7(1); O(35)–Cu(2)–N(26), 122.6(1); N(15)–Cu(2)–N(26), 80.6(1); Cu(2)–O(35)–C(36), 127.3(2).

<sup>†</sup> The analytical data for (**1**) and (**2**) (C, H, N, Cu) were satisfactory.


<sup>‡</sup> Crystal data for (**1**):  $\text{C}_{26}\text{H}_{25}\text{N}_4\text{O}_{10}\text{ClCu}_2$ ,  $M = 716.05$ , triclinic,  $P\bar{1}$ ,  $a = 11.973(2)$ ,  $b = 18.576(4)$ ,  $c = 7.745(1)$  Å,  $\alpha = 103.96(1)$ ,  $\beta = 65.92(1)$ ;  $\gamma = 119.58(1)^\circ$ ,  $U = 1365.95$  Å<sup>3</sup>,  $Z = 2$ ,  $D_c = 1.741$  g cm<sup>-3</sup>,  $\lambda = 0.71069$  Å,  $T = -145^\circ\text{C}$ ,  $6^\circ \leq 2\theta \leq 45^\circ$ ,  $R(R_w) = 2.57(2.91)\%$  for 3271 unique reflections with  $F > 2.33\sigma(F)$ . For (**2**):  $\text{C}_{22}\text{H}_{22}\text{N}_4\text{O}_{12}\text{Cl}_2\text{Cu}_2$ ,  $M = 732.43$ , triclinic,  $P\bar{1}$ ,  $a = 11.286(2)$ ,  $b = 16.414(4)$ ,  $c = 8.047(2)$  Å,  $\alpha = 97.31(1)$ ,  $\beta = 103.78(1)$ ,  $\gamma = 72.59(1)^\circ$ ,  $U = 1378.77$  Å<sup>3</sup>,  $Z = 2$ ,  $D_c = 1.764$  g cm<sup>-3</sup>,  $\lambda = 0.71069$  Å,  $T = -160^\circ\text{C}$ ,  $6^\circ \leq 2\theta \leq 45^\circ$ ,  $R(R_w) = 4.67(4.86)\%$  for 3079 unique reflections with  $F > 3.0\sigma(F)$ . Both structures were solved by MULTAN and Fourier techniques, and refined by full-matrix least-squares. All non-hydrogen atoms were refined anisotropically. Essentially all hydrogen atoms were located in difference Fourier maps (including those of bridging OH<sup>-</sup> and H<sub>2</sub>O groups) and all refined in the final cycles with isotropic thermal parameters. Atomic co-ordinates, bond lengths and angles, and thermal parameters, have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.



**Figure 2.** The structure of the cation of complex (2); OH<sup>-</sup> and H<sub>2</sub>O hydrogen atoms depicted as spheres of arbitrary size. Selected distances (Å) and angles (deg). Cu(1)···Cu(2), 3.035(2); Cu(1)–O(3), 1.927(4); Cu(1)–O(4), 2.405(4); Cu(1)–O(7), 1.931(4); Cu(1)–N(9), 2.006(5); Cu(1)–N(16), 2.008(5); Cu(2)–O(3), 1.930(4); Cu(2)–O(4), 2.379(4); Cu(2)–O(5), 1.944(4); Cu(2)–N(21), 2.010(5); Cu(2)–N(28), 2.009(4); C(6)–O(7), 1.267(7); C(6)–O(5), 1.264(7); Cu(1)–O(3)–Cu(2), 103.8(2); Cu(1)–O(4)–Cu(2), 78.7(1); O(3)–Cu(1)–O(4), 80.3(1); O(3)–Cu(1)–O(7), 95.2(2); O(3)–Cu(1)–N(9), 175.0(2); O(3)–Cu(1)–N(16), 93.9(2); O(4)–Cu(1)–O(7), 98.2(2); O(4)–Cu(1)–N(9), 100.5(2); O(4)–Cu(1)–N(16), 92.9(2); O(7)–Cu(1)–N(9), 89.6(2); O(7)–Cu(1)–N(16), 166.7(2); N(9)–Cu(1)–N(16), 81.2(2); O(3)–Cu(2)–O(4), 80.9(2); O(3)–Cu(2)–O(5), 94.2(2); O(3)–Cu(2)–N(21), 176.3(2); O(3)–Cu(2)–N(28), 95.4(2); O(4)–Cu(2)–O(5), 98.7(2); O(4)–Cu(2)–N(21), 99.4(2); O(4)–Cu(2)–N(28), 98.2(2); O(5)–Cu(2)–N(21), 89.4(2); O(5)–Cu(2)–N(28), 161.6(2); N(21)–Cu(2)–N(28), 80.9(2); Cu(2)–O(5)–C(6), 129.6(4); Cu(1)–O(7)–C(6), 129.4(4).

The structure of complex (2) again consists of a triply-bridged pair of five-coordinate Cu atoms, but only one of the bridging ligands is an AcO<sup>-</sup>, the others being an OH<sup>-</sup> and a H<sub>2</sub>O molecule. As a result of more mono-atomic bridges, the Cu···Cu separation in (2) [3.035(2) Å] is shorter than that in (1) [3.392(1) Å]. The metal co-ordination geometries are square pyramidal with the Cu atoms lying in the basal planes<sup>§</sup> and the H<sub>2</sub>O molecule [O(4)] occupying the apical positions for both metals; the apical Cu(1)–O(4) and Cu(2)–O(4) distances are both quite long [2.405(4) and 2.379(4) Å, respectively] as expected. We are unaware of another structurally-characterized example of a triply-bridged dinuclear Cu complex possessing both OH<sup>-</sup> and H<sub>2</sub>O bridging groups.<sup>6,7</sup>

The only difference in the synthetic procedures to (1) and (2) is the identity of the solvents, complex (2) forming in the predominantly aqueous medium. We thus reasoned that (1) might convert to (2) in H<sub>2</sub>O, and recrystallization of (1) from hot water does indeed yield (2), in essentially quantitative yield if the required extra ClO<sub>4</sub><sup>-</sup> is added [equation (1)].



Variable temperature magnetic susceptibility data were collected on powdered samples of (1) and (2) in the

temperature range 5–300 K. The data were least-squares fit to the Bleaney–Bowers equation<sup>8</sup> replacing  $T$  with  $(T - \theta)$  where  $\theta$  is the Weiss constant. For complex (1), the effective magnetic moment,  $\mu_{\text{eff}}$ , per Cu<sub>2</sub> decreases gradually from 2.96  $\mu_{\text{B}}$  at 295.0 K to a minimum of 2.69  $\mu_{\text{B}}$  at 60.0 K and then rises to 2.87  $\mu_{\text{B}}$  at 5.0 K. The behaviour is clearly characteristic of intramolecular ferromagnetic coupling, and the fitting parameters were  $J = +3.6 \text{ cm}^{-1}$ ,  $g = 2.1$  and  $\theta = 0.20 \text{ K}$ .

Complex (2) is also ferromagnetically-coupled:  $\mu_{\text{eff}}/\text{Cu}_2$  rises from 2.64  $\mu_{\text{B}}$  at 200 K to 2.93  $\mu_{\text{B}}$  at 50.0 K and then is relatively constant with a value of 2.95  $\mu_{\text{B}}$  at 5.0 K. However, the low-temperature plateau could not be modelled satisfactorily, and we believe the cause to be weak intermolecular antiferromagnetic interactions mediated by the hydrogen bonds between the bridging OH<sup>-</sup> and H<sub>2</sub>O groups and the ClO<sub>4</sub><sup>-</sup> anions which give one-dimensional H-bonded chains.<sup>¶</sup> Therefore, only the higher temperature data were fit to provide an estimate of  $J$ , and the fitting parameters were  $J = +19.3 \text{ cm}^{-1}$ ,  $g = 2.1$  and  $\theta = 4.5 \text{ K}$ . Although antiferromagnetically-coupled Cu dimers are by far the more common, ferromagnetic coupling is frequently seen and the values obtained for (1) and (2) fall in the typical range observed.<sup>9</sup>

This work was supported by the National Science Foundation.

Received, 23rd January 1990; Com. 0/00362J

## References

- G. Christou, *Acc. Chem. Res.*, 1989, **22**, 328, and references cited therein.
- J. B. Vincent, J. C. Huffman, G. Christou, Q. Li, M. A. Nanny, D. N. Hendrickson, R. H. Fong, and R. H. Fish, *J. Am. Chem. Soc.*, 1988, **110**, 6898.
- R. C. Mehrotra and R. Bohra, 'Metal Carboxylates,' Academic Press, London, 1983.
- A. M. Greenaway, C. J. O'Connor, J. W. Overman, and E. Sinn, *Inorg. Chem.*, 1981, **20**, 1508; S. J. Brown, X. Tao, D. W. Stephan, and P. Mascharak, *Inorg. Chem.*, 1986, **25**, 3377; W. B. Tolman, A. Bino, and S. J. Lippard, *J. Am. Chem. Soc.*, 1989, **111**, 8522.
- B. Chiari, O. Piovesana, T. Tarantelli, and P. F. Zanazzi, *Inorg. Chem.*, 1985, **24**, 4615; R. Österberg, B. Sjöberg, and R. Soderquist, *J. Chem. Soc., Chem. Commun.*, 1972, 983; B. Chiari, W. E. Hatfield, O. Piovesana, T. Tarantelli, L. W. ter Haar, and P. F. Zanazzi, *Inorg. Chem.*, 1983, **22**, 1468; J.-P. Costes, F. Dahan, and J.-P. Laurent, *Inorg. Chem.*, 1985, **24**, 1018.
- The closest precedent is [Cu<sub>2</sub>(OH)(MeOH)(OAc)(L–L)]<sup>2+</sup> (L–L = a binucleating tetradentate ligand) but the Cu···O (MeOH) distances are much longer [2.59(1) Å] and the authors prefer to describe the complex as an inclusion compound rather than MeOH-bridged: W. B. Tolman, R. L. Rardin, and S. J. Lippard, *J. Am. Chem. Soc.*, 1989, **111**, 4532.
- One dinuclear Cu<sup>II</sup> complex possessing only one OH<sup>-</sup> and one H<sub>2</sub>O bridge is known: M. G. B. Drew, J. Nelson, F. Esho, V. McKee, and S. M. Nelson, *J. Chem. Soc., Dalton Trans.*, 1982, 1837.
- B. Bleaney and K. D. Bowers, *Proc. R. Soc., London, A*, 1952, **214**, 451.
- For some recent reviews, see: M. Melnik, *Coord. Chem. Rev.*, 1982, **42**, 259; 'Magneto-Structural Correlations in Exchange Coupled Systems,' eds. R. D. Willett, D. Gatteschi, and O. Kahn, Reidel, Dordrecht, 1985; M. Kato and Y. Muto, *Coord. Chem. Rev.*, 1988, **92**, 45; O. Kahn, *Inorg. Chim. Acta*, 1982, **62**, 3.

<sup>§</sup> The largest deviation from the least-squares plane Cu(1), O(3), O(7), N(9), N(16) is 0.093 Å by N(16), and from the plane Cu(2), O(3), O(5), N(21), N(28) it is 0.159 Å by N(28). The dihedral angle between these planes is 61.7°.

<sup>¶</sup> The one-dimensional chain of Cu<sub>2</sub> cations is formed by an O atom of one ClO<sub>4</sub><sup>-</sup> being H-bonded both to the OH<sup>-</sup> hydrogen of one Cu<sub>2</sub> cation and one of the H<sub>2</sub>O hydrogens of an adjacent Cu<sub>2</sub> cation. The other ClO<sub>4</sub><sup>-</sup> is H-bonded to the second H<sub>2</sub>O hydrogen but does not bridge to a neighbouring Cu<sub>2</sub> cation.